

National Dose Banding Table – Single Container

Strength of raw material (after reconstitution if required)	25 mg/mL	Example drug(s)	Fludarabine Methotrexate (Std Dose)
--	----------	-----------------	--

See table usage notes below regarding 'single container' and 'multiple syringe' tables.

Master Bands and Ranges

This table is intended to be in a format useful for electronic prescribing systems.

Use To (A) if your system will round UP for doses on the step between two bands.

Use To (B) if your system will round DOWN for doses on the step between two bands.

Band Range (mg)			Band Dose (mg)	Variance (percent)	
From \geq	To (A) <	To (B) \leq		Below	Above
13.73	15.23	15.22	14.5	6	-5
15.23	16.98	16.97	16	5	-6
16.98	18.98	18.97	18	6	-5
18.98	21.23	21.22	20	5	-6
21.23	23.73	23.72	22.5	6	-5
23.73	26.23	26.22	25	5	-5
26.23	28.73	28.72	27.5	5	-4
28.73	31.23	31.22	30	4	-4
31.23	33.73	33.72	32.5	4	-4
33.73	36.23	36.22	35	4	-3
36.23	38.73	38.72	37.5	4	-3
38.73	41.23	41.22	40	3	-3
41.23	43.73	43.72	42.5	3	-3
43.73	47.43	47.42	45	3	-5
47.43	52.45	52.44	50	5	-5
52.45	57.45	57.44	55	5	-4
57.45	62.45	62.44	60	4	-4
62.45	67.45	67.44	65	4	-4
67.45	72.45	72.44	70	4	-3
72.45	77.45	77.44	75	4	-3
77.45	82.45	82.44	80	3	-3
82.45	87.48	87.47	85	3	-3
87.48	94.88	94.87	90	3	-5
94.88	104.88	104.87	100	5	-5
104.88	114.90	114.89	110	5	-4
114.90	127.28	127.27	120	4	-6
127.28	142.30	142.29	135	6	-5
142.30	157.33	157.32	150	5	-5
157.33	174.73	174.72	165	5	-6
174.73	194.75	194.74	185	6	-5
194.75	214.78	214.77	205	5	-5
214.78	237.18	237.17	225	5	-5
237.18	262.20	262.19	250	5	-5
262.20	287.23	287.22	275	5	-4
287.23	312.25	312.24	300	4	-4



Drug- and Dose-specific Tables

These tables are intended to assist with non-electronic systems where protocol-specific information is required. The range of the tables has been extended beyond 'normal' use, especially in the lower ranges, to allow for up to a 50% dose reduction to be used in patients who are $1.3m^2$. Doses which are normally capped are highlighted in grey.

Fludarabine

BSA	Choose nearest for dose reductions							1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
25 mg/m ²	11.5	13	14.5	16	18	20	22.5	25	27.5	30	32.5	35	37.5	40	42.5	45	50	50	55	55	60	60	65	65	70	70	75	75
30 mg/m ²	14.5	16	18	20	22.5	25	27.5	30	32.5	35	40	42.5	45	50	50	55	55	60	65	65	70	70	75	80	80	85	85	90
40 mg/m ²	22.5	25	27.5	30	32.5	35	37.5	40	45	50	50	55	60	65	70	70	75	80	85	90	90	100	100	100	110	110	120	120
50 mg/m ²	30	32.5	35	37.5	40	42.5	45	50	55	60	65	70	75	80	85	90	100	100	110	110	120	120	120	135	135	135	150	150

Methotrexate (Standard Dose)

BSA	Choose nearest for dose reductions							1.0	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0
30 mg/m ²	14.5	16	18	20	22.5	25	27.5	30	32.5	35	40	42.5	45	50	50	50	50	50	50	50	50	50	50	50	50	50	50	50
40 mg/m ²	22.5	25	27.5	30	32.5	35	37.5	40	45	50	50	55	60	65	70	70	75	80	85	90	90	100	100	100	110	110	120	120
100 mg/m ²	60	65	70	75	80	85	90	100	110	120	135	135	150	165	165	185	185	205	205	225	225	250	250	250	275	275	300	300

Note: 30mg/m² used in sarcoma and capped at 50mg.

Table usage notes: 'Single container' tables are for products which are able to be supplied as a single unit, such as an infusion bag, one syringe, or as an infusor device (pump). 'Multiple syringe' (e.g. 'pick-and-mix') tables are for products which are supplied in combinations of syringes to make up a single dose. Some drugs may have both a single container table, and a multiple syringe table. Doses may vary between the two tables due to rounding requirements.

